

Perivasculo-Ductal Connective Tissue in the Parotid Gland of Wistar Rats Exposed to Low Frequency Noise.


Pedro Oliveira, PhD¹, José Martins dos Santos, PhD¹, José João Mendes, DDS¹, Mariana Alves-Pereira, MSc², Nuno Castelo Branco, MD³ Egas Moniz School of Health Sciences, Qta da Granja, Monte da Caparica, 2829-511 Portugal. Phone: +351-964588445, email: pedromaoliveira@hotmail.com. ²ERISA-Universidade Lusófona, R. do Telhal aos Olivais, 1900 Lisbon, Portugal, phone:+351218621060, email: m.alvespereira@gmail.com. ³Center for Human Performance, R. Prof. Dias Amado, No. 5, 1B, 1600 Lisbon, Portugal, phone: +351-93-7584261, email: vibroacoustic.disease@gmail.com.

BACKGROUND: In vibroacoustic disease (VAD), fibrosis occurs in the absence of inflammatory signs, and is thought to be a protective response of tissues and organs to the presence of the corresponding agent of disease: low frequency noise (LFN, ≤ 500 Hz). In the parotid gland, the perivasculo-ductal connective tissue (PVDCT) follows the arteries, veins, and ductal tree, and is thought to function as a mechanical stabilizer of the glandular tissue [1].

PURPOSE: To quantify the proliferation of PVDCT in LFN-exposed rats.

METHODS: Sixty Wistar rats were equally divided into 6 groups, one kept in silence (gC), and the remaining 5 exposed to continuous LFN for different amounts of time: g1- 168h (1wk); g2- 504 hrs (3wks); g3- 840 hrs (5wks); g4- 1512 hrs (9wks); and g5- 2184 hrs (13 weeks). After exposure, parotid glands were removed and the PVDCT perimeter was measured in all groups.

RESULTS: See Chart. The global trend is an increase in the average PVDCT areas, that develops linearly and significantly with LFN exposure time ($p < 0001$).



CONCLUSIONS: It has been suggested that the biological response to LFN exposure is associated with the necessity of maintaining structural integrity [2]. The parotid gland PVDCT contains two populations of collagen fibrils (one associated with fibroblasts and the other with epithelium), together with fibroblasts and the epithelium of excretory ducts and blood vessels [1]. In the parotid gland, structural reinforcement would be achieved at the cost of increased PVDCT. Hence, these results show that in response to LFN exposure, rat parotid glands increase their PVDCT.

REFERENCES:

1. Hosoyamada Y, Sakai T. The ultrastructure of periductal connective tissue and distinctive populations of collagen fibrils associated with ductal epithelia of exocrine glands. *Arch Histol Cytol*, 66: 407-18, 2003.
2. Alves-Pereira M, Castelo Branco NAA. Vibroacoustic disease: Biological effects of infrasound and low frequency noise explained by mechanotransduction cellular signaling. *Prog Biophy Molec Biol*, 93:256-279, 2007.